

Lecture-25: Poisson processes: Conditional distribution

1 Joint conditional distribution of points in a finite window

Let $\mathcal{X} = \mathbb{R}^d$ be a d -dimensional Euclidean space.

Proposition 1.1. *Let $k \in \mathbb{N}$ be any positive integer. For a Poisson point process $S : \Omega \rightarrow \mathcal{X}^{\mathbb{N}}$ with intensity measure $\Lambda : \mathcal{B}(\mathcal{X}) \rightarrow \mathbb{R}_+$, consider a bounded subset $A \in \mathcal{B}(\mathcal{X})$ and subsets $(A_1, \dots, A_k) \in \mathcal{B}(\mathcal{X})^k$ that partition A . Let $n_1, \dots, n_k \in \mathbb{Z}_+$ such that $n_1 + \dots + n_k = n$, then*

$$P(\{N(A_1) = n_1, \dots, N(A_k) = n_k\} \mid \{N(A) = n\}) = \binom{n}{n_1, \dots, n_k} \prod_{i=1}^k \left(\frac{\Lambda(A_i)}{\Lambda(A)} \right)^{n_i}. \quad (1)$$

Proof. It follows from the definition of joint distribution of $(N(A_1), \dots, N(A_k))$, the fact that $\bigcap_{i=1}^k \{N(A_i) = n_i\} \subseteq \{N(A) = n\}$, and that the intensity measure add over disjoint sets, i.e. $\Lambda(A) = \sum_{i=1}^k \Lambda(A_i)$. \square

Remark 1. Let $S : \Omega \rightarrow \mathcal{X}^{\mathbb{N}}$ be a Poisson point process with intensity measure $\Lambda : \mathcal{B}(\mathcal{X}) \rightarrow \mathbb{R}_+$, and $(A_1, \dots, A_k) \in \mathcal{B}(\mathcal{X})^k$ be disjoint bounded subsets such that $A = \cup_{i=1}^k A_i \in \mathcal{B}(\mathcal{X})$. Since S is a simple point process, each point S_n is unique. Therefore, we can identify the random sequence S of points in \mathcal{X} as a random set of points in \mathcal{X} . It follows that $\{N(A) = n\} = \{|S \cap A| = n\}$.

i. From the disjointness of A_i , we have $N(A) = N(A_1) + \dots + N(A_k)$, and from the linearity of expectations, we get

$$\Lambda(A) = \mathbb{E}N(A) = \sum_{i=1}^k \mathbb{E}N(A_i) = \sum_{i=1}^k \Lambda(A_i).$$

ii. Defining $p_i \triangleq \frac{\Lambda(A_i)}{\Lambda(A)}$, we see that (p_1, \dots, p_k) is a probability distribution. We also observe that

$$p_i = P(\{N(A_i) = 1\} \mid \{N(A) = 1\}) = P(\{|S \cap A_i| = 1\} \mid \{|S \cap A| = 1\}).$$

When $N(A) = 1$, we can call the point of S in A as S_1 WLOG. That is, we let $\{S_1\} = S \cap A$, to observe

$$p_i = P(\{S_1 \in A_i\} \mid \{S_1 \in A\}).$$

Similarly, when $N(A) = n_i$, we call the points of S in A as S_1, \dots, S_{n_i} . That is, we let $\{S_1, \dots, S_{n_i}\} = S \cap A$, to observe

$$\begin{aligned} p_i^{n_i} &= P(\{N(A_i) = n_i\} \mid \{N(A) = n_i\}) = P(\{|S \cap A_i| = n_i\} \mid \{|S \cap A| = n_i\}) \\ &= P(\{\bigcap_{j=1}^{n_i} \{S_j \in A_i\}\} \mid \{\{S_1, \dots, S_{n_i}\} = S \cap A\}) = \prod_{j=1}^{n_i} P(\{S_j \in A_i\} \mid \{S_j \in A\}). \end{aligned}$$

iii. We can rewrite the Equation (1) as a multinomial distribution, where

$$P(\{N(A_1) = n_1, \dots, N(A_k) = n_k\} \mid \{N(A) = n\}) = \binom{n}{n_1, \dots, n_k} p_1^{n_1} \dots p_k^{n_k}.$$

iv. Let $A \in \mathcal{B}(X)$ be a bounded set, $\mathcal{P}(A)$ be the set of all subsets of A . Let $\mathcal{P}_k(A)$ be the collection of k -partitions (E_1, \dots, E_k) of set A . That is,

$$\mathcal{P}_k(A) = \left\{ (E_1, \dots, E_k) \in \mathcal{P}([n])^k : E_i \cap E_j = \emptyset, i \neq j, \cup_{i=1}^k E_i = A \right\}.$$

We define $\mathcal{P}_k(n_1, \dots, n_k)$ to be the collection of all k -partitions (E_1, \dots, E_k) of any finite set $A \subseteq \mathcal{B}(X)$ such that $|E_i| = n_i$ for $i \in [k]$ and $n_1 + \dots + n_k = |A|$. That is,

$$\mathcal{P}_k(A, n_1, \dots, n_k) \triangleq \{(E_1, \dots, E_k) \in \mathcal{P}_k(A) : |E_i| = n_i \text{ for all } i \in [k]\}.$$

Then, the multinomial coefficient accounts for number of partitions of n points into sets with n_1, \dots, n_k points. That is,

$$\binom{n}{n_1, \dots, n_k} = |\mathcal{P}_k([n], n_1, \dots, n_k)|.$$

v. Recall that the event $\{N(A_i) = n_i\} = \{|S \cap A_i| = n_i\}$. Hence, we can write

$$\begin{aligned} P(\cap_{i=1}^k \{|S \cap A_i| = n_i\} \mid \{|S \cap A| = n\}) &= \binom{n}{n_1, \dots, n_k} p_1^{n_1} \dots p_k^{n_k} \\ &= \sum_{(E_1, \dots, E_k) \in \mathcal{P}_k(S \cap A, n_1, \dots, n_k)} \prod_{i=1}^k \prod_{S_j \in E_i} P(\{S_j \in A_i\} \mid \{S_j \in A\}). \end{aligned}$$

vi. We further observe that when $N(A_i) = n_i$ for all $i \in [k]$, then $(S \cap A_1, \dots, S \cap A_k) \in \mathcal{P}_k(S \cap A, n_1, \dots, n_k)$. Therefore, we can re-write the event

$$\cap_{i=1}^k \{N(A_i) = n_i\} = \cap_{i=1}^k \{|S \cap A_i| = n_i\} = \cup_{(E_1, \dots, E_k) \in \mathcal{P}_k(S \cap A, n_1, \dots, n_k)} (\cap_{i=1}^k \{S \cap A_i = E_i\}).$$

That is, we can write the conditional probability

$$\begin{aligned} P(\cap_{i=1}^k \{N(A_i) = n_i\} \mid \{N(A) = n\}) &= \sum_{(E_1, \dots, E_k) \in \mathcal{P}_k(S \cap A, n_1, \dots, n_k)} P(\cap_{i=1}^k \{S \cap A_i = E_i\} \mid \{S \cap A = E\}) \\ &= \sum_{(E_1, \dots, E_k) \in \mathcal{P}_k(S \cap A, n_1, \dots, n_k)} P(\cap_{i=1}^k \cap_{S_j \in E_i} \{S_j \in A_i\} \mid \{S \cap A = E\}). \end{aligned}$$

vii. Let S_1, \dots, S_n be the n points in $E = S \cap A$. Equating the RHS of the above equation term-wise, we obtain that conditioned on each of these points falling inside the window A , the conditional probability of each point falling in partition A_i is independent of all other points and given by p_i . That is, we have

$$P(\cap_{i=1}^k \cap_{S_j \in E_i} \{S_j \in A_i\} \mid \{S \cap A = E\}) = \prod_{i=1}^k \prod_{S_j \in E_i} P(\{S_j \in A_i\} \mid \{S_j \in A\}) = \prod_{i=1}^k p_i^{n_i} = \prod_{i=1}^k \left(\frac{\Lambda(A_i)}{\Lambda(A)} \right)^{n_i}.$$

It means that given n points in the window A , the location of these points are independently and identically distributed in A according to the distribution $\frac{\Lambda(\cdot)}{\Lambda(A)}$.

viii. If the Poisson process is homogeneous, the distribution is uniform over the window A .

ix. For a Poisson process with intensity measure Λ and any bounded set $A \in \mathcal{B}$, we have $N(A)$ a Poisson random variable with parameter $\Lambda(A)$. Given $N(A)$, the location of all the points in $S \cap A$ are *i.i.d.* with density $\frac{\lambda(x)}{\Lambda(A)}$ for all $x \in A$.

Corollary 1.2. For a homogeneous Poisson point process on the half-line with ordered set of points $(S_{(n)} \in \mathbb{R}_+ :$

$n \in \mathbb{N}$), we can write the conditional density of ordered points $(S_{(1)}, \dots, S_{(k)})$ given $N(t) = k$ as ordered statistics of i.i.d. uniformly distributed random variables. Specifically, we have

$$f_{S_{(1)}, \dots, S_{(k)} \mid N(t)=k}(t_1, \dots, t_k) = k! \frac{\mathbb{1}_{\{0 < t_1 \leq \dots \leq t_k \leq t\}}}{t}.$$

Proof. Given $\{N(t) = k\}$, we can denote the points of the Poisson process in $(0, t]$ by S_1, \dots, S_k . From the above remark, we know that S_1, \dots, S_k are i.i.d. uniform in $(0, t]$, conditioned on the number of points $N(t) = k$. Hence, we can write

$$F_{S_1, \dots, S_k \mid N(t)=k}(t_1, \dots, t_k) = P(\cap_{i=1}^k \{S_i \in (0, t_i]\} \mid \{N(t) = k\}) = \prod_{i=1}^k P(\{S_i \in (0, t_i]\} \mid \{S_i \in (0, t]\}) = \prod_{i=1}^k \frac{t_i}{t} \mathbb{1}_{\{0 < t_i \leq t\}}.$$

It follows that for any permutation $\sigma : [k] \rightarrow [k]$, the joint distribution of $(S_{\sigma(1)}, \dots, S_{\sigma(k)})$ is identical to that of (S_1, \dots, S_k) . Further, we observe that the order statistics of $(S_{\sigma(1)}, \dots, S_{\sigma(k)})$ is identical to that of (S_1, \dots, S_k) . Therefore, we can write the following equality for the events

$$\{0 < S_{(1)} \leq \dots \leq S_{(k)} \leq t_k \leq t\} = \cup_{\sigma: [k] \rightarrow [k] \text{ permutation}} \cap_{i=1}^k \{S_{\sigma(i)} \leq t_i\}.$$

The result follows since the number of permutations $\sigma : [k] \rightarrow [k]$ is $k!$. □