

Lecture-02: Review of Linear Algebra and Convex Optimization

1 Review of Linear Algebra

1.1 Vector Space

Definition 1.1 (Vector addition). A set V is set to be equipped with vector addition mapping $+: V \times V \rightarrow V$ defined by $+(v, w) = v + w$ for any two elements $v, w \in V$, if this mapping satisfies the following four axioms.

Associativity: $u + (v + w) = (u + v) + w$; for all $u, v, w \in V$.

Commutativity: $u + v = v + u$; for all $u, v \in V$.

Additive identity: There exists a zero vector ($0 \in V$) s.t. $u + 0 = u$; for all $u \in V$.

Additive inverse: For every $u \in V$, there exists an element $-u \in V$; s.t. $u + (-u) = 0$.

Definition 1.2 (Scalar multiplication). A set V equipped with vector addition $+: V \times V \rightarrow V$ is also equipped with field scalar multiplication mapping $\cdot: \mathbb{F} \times V \rightarrow V$ defined by $\cdot(\alpha, v) = \alpha v \in V$, if this mapping satisfies the following four axioms.

Field compatibility: $a(bu) = (ab)u$; for all $a, b \in \mathbb{F}$ and $u \in V$.

Multiplicative identity: For multiplicative identity element $1 \in \mathbb{F}$, $1u = u$; for all $u \in V$.

Distributivity over vector addition: $\alpha(vu) = \alpha u + \alpha v$; for all $\alpha \in \mathbb{F}$ and $u, v \in V$.

Distributivity over field addition: $(\alpha + \beta)u = \alpha u + \beta u$; for all $\alpha, \beta \in \mathbb{F}$ and $u \in V$.

Definition 1.3 (Vector space). A vector space over the field \mathbb{F} is a set V equipped with vector addition $+: V \times V \rightarrow V$ and scalar multiplication $\cdot: \mathbb{F} \times V \rightarrow V$.

Definition 1.4. A set of vectors $W \subseteq V$ are called linearly independent, if for any nonzero vector $\alpha \in \mathbb{F}^W$ with finite $\sum_w \alpha_w$, we have $\sum_{w \in W} \alpha_w w \neq 0 \in V$.

Definition 1.5. The span of a set of vectors $W \subseteq V$ is defined by $\text{span}(W) \triangleq \{ \sum_{w \in W} \alpha_w w : \alpha \in \mathbb{R}^W, \sum_{w \in W} \alpha_w \text{ finite} \}$.

Definition 1.6. A basis of any vector space V , is a spanning set of linearly independent vectors.

Theorem 1.7. All bases of a vector space V have identical cardinality, and defined to be its dimension.

Example 1.8 (Vector space). Following are some common examples of vector spaces.

1. Euclidean space of N -dimensions, denoted by \mathbb{R}^N .
2. Space of continuous functions over a compact subset $[a, b]$ denoted by $C([a, b])$.
3. Space of random variables defined over probability space (Ω, \mathcal{F}, P) .

1.2 Inner Product Space

A *inner product space* is a vector space equipped with an inner product denoted by $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{R}$ that satisfies the following axioms.

1. **Symmetry:** $\langle x, y \rangle = \langle y, x \rangle$
2. **Linearity:** $\langle \alpha x + \beta y, z \rangle = \alpha \langle x, z \rangle + \beta \langle y, z \rangle$
3. **Definiteness:** $\langle x, x \rangle \geq 0$; $\langle x, x \rangle = 0$ iff $x = 0$

Example 1.9 (inner product spaces). Following are some common examples of inner product spaces.

1. For the vector space $V = \mathbb{R}^N$ of N -dimensional vectors, the inner product is defined as $\langle x, y \rangle \triangleq x^T y = \sum_i^N x_i y_i$.
2. For vector space $V = C(\mathbb{R}^N)$ of continuous functions, the inner product is defined as $\langle f, g \rangle \triangleq \int_{\mathbb{R}^N} (f, g)(t) dt$.
3. For the vector space of random variables, the inner product is defined as $\langle X, Y \rangle \triangleq \mathbb{E}XY$.

1.3 Norms

Norm is a mapping $\|\cdot\| : V \rightarrow \mathbb{R}_+$ that satisfy the following axioms.

1. **Definiteness:** $\|v\| = 0$ iff $v = 0$
2. **Homogeneity:** $\|\alpha v\| = |\alpha| \|v\|$
3. **Triangle inequality:** $\|v + w\| \leq \|v\| + \|w\|$

Example 1.10 (Norms). For a vector space $V = \mathbb{R}^N$ of N dimensional vectors, we can define the p -norm for $p > 1$ as $\|x\|_p \triangleq \left(\sum_{i=1}^N |x_i|^p \right)^{\frac{1}{p}}$ for all $x \in \mathbb{R}^N$. For $p = 1$, we have $\|x\|_1 = \sum_{i=1}^N |x_i|$. For $p = \infty$, we have $\|x\|_\infty = \max_i |x_i|$. For $p = 2$, the norm is Euclidean norm.

Proposition 1.11 (Holder's Inequality). Let $p, q \geq 1$ be conjugate, i.e. $\frac{1}{p} + \frac{1}{q} = 1$ Then,

$$|\langle x, y \rangle| \leq \|x\|_p \|y\|_q \text{ for all } x, y \in \mathbb{R}^N.$$

Proof. The Holder's inequality is trivially true if $x = 0$ or $y = 0$. Hence, we assume that $\|x\| \|y\| > 0$, and let $a \triangleq \frac{|x_i|}{\|x\|_p}$ and $b \triangleq \frac{|y_i|}{\|y\|_q}$. We will use the Young's inequality $\frac{1}{p}a^p + \frac{1}{q}b^q \geq ab$ for all $a, b > 0$, that implies that

$$\frac{|x_i|^p}{p \|x\|_p^p} + \frac{|y_i|^q}{q \|y\|_q^q} \geq \frac{|x_i| |y_i|}{\|x\|_p \|y\|_q}, \text{ for all } i \in [N].$$

Since $|\langle x, y \rangle| \leq \sum_{i=1}^N |x_i| |y_i|$, we get the result by summing both sides over $i \in [N]$ in the above inequality.

2 Review of Convex Optimization

Let $\mathcal{X} \subseteq \mathbb{R}^N$ for $N \geq 1$ and $f : \mathcal{X} \rightarrow \mathbb{R}$ be a smooth function.

Definition 2.1 (Gradient). The gradient of function f at point $x \in \mathcal{X}$ is defined as the column vector $\nabla f(x) \in \mathbb{R}^N$, where the i th entry is defined as $\nabla f_i(x) \triangleq \frac{\partial f}{\partial x_i}(x)$ for all $i \in [N]$.

Definition 2.2 (Hessian). The Hessian of function f at point $x \in \mathcal{X}$ is denoted by the matrix $\nabla^2 f(x) \in \mathbb{R}^{N \times N}$, where the (i, j) th entry is defined as $\nabla^2 f_{i,j}(x) \triangleq \frac{\partial^2 f}{\partial x_i \partial x_j}(x)$ for all $i, j \in [N]$.

Remark 1. Let $f : \mathbb{R}^N \rightarrow \mathbb{R}$ be a smooth function over N -dimensional reals. Then, we can write its Taylor series expansion around the neighborhood of $x \in \mathbb{R}^N$, in terms of the gradient vector $\nabla f(x) \in \mathbb{R}^N$ and the Hessian matrix $\nabla^2 f(x) \in \mathbb{R}^{N \times N}$, as

$$f(y) = f(x) + \langle \nabla f(x), (y-x) \rangle + \frac{1}{2} \langle (y-x), \nabla^2 f(x) (y-x) \rangle + o(\|y-x\|_2^2). \quad (1)$$

Definition 2.3 (Stationary Point). A point $x \in \mathcal{X}$ is called a stationary point, if f attains a local extremum at x .

Remark 2. If $f : \mathcal{X} \rightarrow \mathbb{R}$ is smooth, then $\nabla f(x) = 0$ at a stationary point $x \in \mathcal{X}$.

2.1 Convexity

Definition 2.4 (Convex Set). A set \mathcal{X} is called convex if for all $x, y \in \mathcal{X}$ and $\alpha \in [0, 1]$, the convex combination $\alpha x + \bar{\alpha} y \in \mathcal{X}$ where $\bar{\alpha} \triangleq (1 - \alpha)$.

Definition 2.5 (Convex Hull). A convex hull of a set A is the smallest convex set including A , i.e. $\text{conv}(A) \triangleq \{ \sum_{x \in A} \alpha_x x : 0 \leq \alpha_x \leq 1, \sum_{x \in A} \alpha_x = 1 \}$.

Definition 2.6. Let $\mathcal{X} \subseteq \mathbb{R}^N$. For a function $f : \mathcal{X} \rightarrow \mathbb{R}$, we define its epigraph as

$$\text{Epi}(f) \triangleq \{(x, y) \in \mathcal{X} \times \mathbb{R} : y \geq f(x)\}.$$

Definition 2.7. A function $f : \mathcal{X} \rightarrow \mathbb{R}$ is convex if the associated domain \mathcal{X} and epigraph $\text{Epi}(f)$ are convex sets.

Theorem 2.8. Let $\mathcal{X} \subset \mathbb{R}^N$ be a convex set. Then the following are equivalent statements.

1. $f : \mathcal{X} \rightarrow \mathbb{R}$ is a convex function.

2. For all $\alpha \in [0, 1]$, we have $f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y)$.
3. For differentiable f , we have $f(y) - f(x) \geq \langle \nabla f(x), y - x \rangle$ for all $x, y \in \mathcal{X}$.
4. For twice differentiable f , we have $\nabla^2 f \succeq 0$, i.e. $\nabla^2 f$ is a positive semi-definite matrix.

Proof. For convex set $\mathcal{X} \subseteq \mathbb{R}^N$ and a function $f : \mathcal{X} \rightarrow \mathbb{R}$, we will show that statement 1 implies statement 2, which implies statement 3, which implies statement 4, which implies statement 1.

- 1 \implies 2: Let $(x, f(x)), (y, f(y)) \in \text{Epi}(f)$ for $x, y \in \mathcal{X}$. Let $\alpha \in [0, 1]$, then from the convexity of \mathcal{X} , we have $\alpha x + \bar{\alpha}y \in \mathcal{X}$. Further from the convexity of $\text{Epi}(f)$, we have $(\alpha x + \bar{\alpha}y, \alpha f(x) + \bar{\alpha}f(y)) \in \text{Epi}(f)$. That is, $\alpha f(x) + \bar{\alpha}f(y) \geq f(\alpha x + \bar{\alpha}y)$.
- 2 \implies 3: Recall that $\alpha x + \bar{\alpha}y = x + \bar{\alpha}(y - x)$. From statement 2, we have $f(y) - f(x) \geq \frac{f(\alpha x + \bar{\alpha}y) - f(x)}{\bar{\alpha}}$. Taking $\bar{\alpha} \rightarrow 0$, we observe that the right hand side is equal to $\langle \nabla f(x), y - x \rangle$.
- 3 \implies 4: From (1) and statement 3, it follows that for any $x, y \in \mathcal{X}$ $f(y) - f(x) - \langle \nabla f(x), y - x \rangle = \frac{1}{2}(y - x)^T \nabla^2 f(x)(y - x) + o(\|y - x\|_2^2) \geq 0$.
- 4 \implies 1: Let $\alpha \in [0, 1]$. Then, it suffices to show that $\alpha f(x_1) + \bar{\alpha}f(x_2) \geq f(\alpha x_1 + \bar{\alpha}x_2)$. From the Taylor expansion of f in the neighborhood of x_2 , we get

$$\alpha(f(x_1) - f(x_2)) = \alpha \langle \nabla f(x_2), x_1 - x_2 \rangle + \frac{\alpha}{2} \langle (x_1 - x_2), \nabla^2 f(x_2)(x_1 - x_2) \rangle + o(\|x_1 - x_2\|_2^2).$$

Similarly, we write the Taylor expansion of f in the neighborhood of x_2 , to get

$$f(\alpha x_1 + \bar{\alpha}x_2) - f(x_2) = \alpha \langle \nabla f(x_2), x_1 - x_2 \rangle + \frac{\alpha^2}{2} \langle (x_1 - x_2), \nabla^2 f(x_2)(x_1 - x_2) \rangle + o(\|x_1 - x_2\|_2^2).$$

Taking the difference, we get $\alpha(f(x_1) - f(x_2)) \geq f(\alpha x_1 + \bar{\alpha}x_2) - f(x_2)$. □

Example 2.9 (Convex Function). Following functions $f : \mathbb{R}^N \rightarrow \mathbb{R}$ are convex.

1. Linear Function: $f(x) = \langle w, x \rangle$ for $w \in \mathbb{R}^N$.
2. Quadratic Function: $f(x) = x^T A x$ for a positive semi definite matrix $A \in \mathbb{R}^{N \times N}$.
3. Abs Maximum: $f(x) = \max \{|x_i| : i \in [N]\} = \|x\|_\infty$.

Lemma 2.10 (Composition of functions). We define a composition function $f = h \circ g$ for functions $h : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R}^N \rightarrow \mathbb{R}$, by defining $f(x) \triangleq h(g(x))$ for all $x \in \mathbb{R}^N$. Then, the following statements are true.

1. If h is convex and nondecreasing and g is convex, then f is convex.
2. If h is convex and nonincreasing and g is concave, then f is convex.
3. If h is concave and nondecreasing and g is concave, then f is concave.
4. If h is concave and nonincreasing and g is convex, then f is concave.

Proof. We will use the property that a function f is convex iff $\text{dom}(f)$ is convex and $f(\alpha x + \bar{\alpha}y) \leq \alpha f(x) + \bar{\alpha}f(y)$ for all $\alpha \in [0, 1]$. Recall that \mathbb{R}^N is convex for all $N \geq 1$. We will only show the first statement, and rest follow the same steps. Let $x, y \in \mathbb{R}^N$ and $\alpha \in [0, 1]$. From the convexity of g , we get $g(\alpha x + \bar{\alpha}y) \leq \alpha g(x) + \bar{\alpha}g(y)$. From the nondecreasing property of h , we get $h(g(\alpha x + \bar{\alpha}y)) \leq h(\alpha g(x) + \bar{\alpha}g(y))$. From the convexity of h , we get $h(\alpha g(x) + \bar{\alpha}g(y)) \leq \alpha h(g(x)) + \bar{\alpha}h(g(y))$. □

Theorem 2.11 (Jensen's Inequality). Let $X : \Omega \rightarrow \mathcal{X} \subseteq \mathbb{R}^N$ be a random vector with finite marginal means, and $f : \mathcal{X} \rightarrow \mathbb{R}$ be a convex function. Then the mean $\mathbb{E}[X] \in \mathcal{X}$, the mean $\mathbb{E}[f(X)]$ is finite, and $f(\mathbb{E}[X]) \leq \mathbb{E}[f(X)]$.

Proof. We will show this for simple random vector $X : \Omega \rightarrow \{x_1, \dots, x_m\} \subseteq \mathcal{X}$, such that $\alpha_i \triangleq P\{X = x_i\}$ for all $i \in [m]$. Then, the mean $\mathbb{E}X = \sum_{i=1}^m \alpha_i x_i \in \mathcal{X}$ from the convexity of \mathcal{X} , and $\mathbb{E}f(X) = \sum_{i=1}^m \alpha_i f(x_i)$ is finite. Further, from the convexity of f , we get $f\left(\sum_{i=1}^m \alpha_i x_i\right) \leq \sum_{i=1}^m \alpha_i f(x_i)$. □

Corollary 2.12 (Young's inequality). Let $p, q \geq 1$ be the conjugate pair such that $1/p + 1/q = 1$. Then, $ab \leq \frac{a^p}{p} + \frac{b^q}{q}$.

Proof. Take a random variable $X : \Omega \rightarrow \{a^p, b^q\}$ with probability mass function $P_X(a^p) = \frac{1}{p}$ and $P_X(b^q) = \frac{1}{q}$. Then, from the concavity of log

$$\ln \left(\frac{1}{p}a^p + \frac{1}{q}b^q \right) = \ln \mathbb{E}X \geq \mathbb{E} \ln X = \frac{1}{p} \ln a^p + \frac{1}{q} \ln b^q = \ln ab.$$

Since $\ln(\cdot)$ is an increasing function, the above inequality implies the result. \square

2.2 Constrained Optimization

Problem 2.13 (primal problem). Consider a cost function $f : \mathbb{R}^N \rightarrow \mathbb{R}$ and a constraint function $g : \mathbb{R}^N \rightarrow \mathbb{R}^m$. The **primal problem** is $p^* \triangleq \inf \{f(x) : x \in \mathcal{X}\}$, where the constraint set is

$$\mathcal{X} \triangleq \bigcap_{i=1}^m \{x \in \mathbb{R}^N : g_i(x) \leq 0\}. \quad (2)$$

Definition 2.14 (Lagrangian). For the Problem 2.13, we define an associated Lagrangian function $\mathcal{L} : \mathbb{R}^N \times \mathbb{R}_+^m \rightarrow \mathbb{R}$ for Lagrange or dual variables $\alpha \in \mathbb{R}_+^m$ and primal variables $x \in \mathbb{R}^N$, as

$$\mathcal{L}(x, \alpha) \triangleq f(x) + \langle \alpha, g(x) \rangle. \quad (3)$$

Definition 2.15 (Dual function). The dual function $F : \mathbb{R}_+^m \rightarrow \mathbb{R}$ associated with the Problem 2.13 is defined for dual variables $\alpha \in \mathbb{R}_+^m$ as

$$F(\alpha) \triangleq \inf \{ \mathcal{L}(x, \alpha) : x \in \mathbb{R}^N \}. \quad (4)$$

Theorem 2.16. The following are true for the dual function $F : \mathbb{R}_+^m \rightarrow \mathbb{R}$ defined in (4) for the Problem 2.13.

1. F is concave in $\alpha \in \mathbb{R}_+^m$.
2. $F(\alpha) \leq \mathcal{L}(x, \alpha)$ for all $\alpha \in \mathbb{R}_+^m$ and $x \in \mathbb{R}^N$.
3. $F(\alpha) \leq p^*$ for all $\alpha \in \mathbb{R}_+^m$.

Proof. Recall that $\mathcal{L}(\alpha) = f(x) + \langle \alpha, g(x) \rangle$ is a linear function of $\alpha \in \mathbb{R}_+^m$, and $F(\alpha) = \inf_x \mathcal{L}(x, \alpha)$.

1. Let $\beta \in [0, 1]$ and $\alpha_1, \alpha_2 \in \mathbb{R}_+^m$ and $x \in \mathcal{X}$. It follows from the linearity of Lagrangian in α that

$$F(\beta\alpha_1 + \bar{\beta}\alpha_2) = \inf_x \left[\beta \mathcal{L}(x, \alpha_1) + \bar{\beta} \mathcal{L}(x, \alpha_2) \right] \geq \beta \inf_x \mathcal{L}(x, \alpha_1) + \bar{\beta} \inf_x \mathcal{L}(x, \alpha_2) = \beta F(\alpha_1) + \bar{\beta} F(\alpha_2).$$

2. From the definition of F , it follows that $F(\alpha) \leq \mathcal{L}(x, \alpha)$ for all $x \in \mathbb{R}^N$.
3. Recall that $g_i(x) \leq 0$ for all $x \in \mathcal{X}$, and hence $\langle \alpha, g(x) \rangle \leq 0$ for all $x \in \mathcal{X}$. Therefore, $F(\alpha) \leq f(x)$ for all $x \in \mathcal{X}$, and hence the result follows. \square

Problem 2.17. Dual problem The dual problem associated with primal problem defined in Problem 2.13 is

$$d^* \triangleq \max \{ F(\alpha) : \alpha \in \mathbb{R}_+^m \}.$$

Remark 3. From the properties of dual function $F : \mathbb{R}_+^m \rightarrow \mathbb{R}$ in Theorem 2.16, we obtain that F is concave in $\alpha \in \mathbb{R}_+^m$. Since \mathbb{R}_+^m is a convex set, it follows that the dual problem is convex. We further observe that the optimal value of dual problem $d^* \leq p^*$. The difference of optimal values ($p^* - d^*$) is called the **duality gap**. For a primal problem, the **strong duality** holds if the duality gap is zero, or $d^* = p^*$.

2.3 Convex constrained optimization

Definition 2.18 (Saddle point). For a Lagrangian $\mathcal{L} : \mathbb{R}^N \times \mathbb{R}_+^m \rightarrow \mathbb{R}$, a saddle point (x^*, α^*) satisfies

$$\sup_{\alpha \in \mathbb{R}_+^m} \mathcal{L}(x^*, \alpha) \leq \mathcal{L}(x^*, \alpha^*) \leq \inf_{x \in \mathbb{R}^N} \mathcal{L}(x, \alpha^*).$$

Theorem 2.19 (Sufficient condition). For the primal problem defined in Problem 2.13, if (x^*, α^*) is a saddle point of the associated Lagrangian \mathcal{L} , then $x^* \in \mathcal{X}$ and $p^* = f(x^*) = F(\alpha^*)$.

Proof. Let (x^*, α^*) be the saddle point of the Lagrangian \mathcal{L} associated with the Problem 2.13. From the definition of dual function F , we get that $\mathcal{L}(x^*, \alpha^*) \leq F(\alpha^*) \leq \mathcal{L}(x^*, \alpha^*)$. It follows that $F(\alpha^*) = \mathcal{L}(x^*, \alpha^*)$.

Recall that $\mathcal{L}(x^*, \alpha) = f(x^*) + \langle \alpha, g(x^*) \rangle$. We assume that there exists an $i \in [m]$ such that $g_i(x) > 0$, then we can take α_i large enough so that $\mathcal{L}(x^*, \alpha) \geq \mathcal{L}(x^*, \alpha^*)$. This contradicts the saddle point condition, and hence $x^* \in \mathcal{X}$. Therefore $\langle \alpha, g(x^*) \rangle \leq 0$ for all $\alpha \in \mathbb{R}_+^m$. This implies that $\langle \alpha^*, g(x^*) \rangle = 0$ and hence $p^* = f(x^*) = F(\alpha^*)$. \square

Definition 2.20 (Strong constraint qualification). The strong constraint qualification or **Slater's condition** is defined as the existence of a point $x \in \mathcal{X}^\circ$ such that $g_i(x) < 0$ for all $i \in [m]$.

Theorem 2.21 (Strong necessary condition). *Let the cost function f and constraints g_i for $i \in [m]$ be convex functions, such that the Slater's condition holds, and x^* be the solution of the Problem 2.13. Then, there exists $\alpha^* \in \mathbb{R}_+^m$ such that (x^*, α^*) is a saddle point of the associated Lagrangian \mathcal{L} .*

Definition 2.22 (Weak constraint qualification). The weak constraint qualification or **weak Slater's condition** is defined as the existence of a point $x \in \mathcal{X}^\circ$ such that for each $i \in [m]$ either $g_i(x) < 0$ or $g_i(x) = 0$ and g_i affine.

Theorem 2.23 (Weak necessary condition). *Let the cost function f and constraints g_i for $i \in [m]$ be convex differentiable functions, such that the weak Slater's condition holds, and x^* be the solution of the Problem 2.13. Then, there exists $\alpha^* \in \mathbb{R}_+^m$ such that (x^*, α^*) is a saddle point of the associated Lagrangian \mathcal{L} .*

Remark 4. The strong duality holds when the primal problem is convex with qualifying constraints.

Theorem 2.24 (Karush-Kuhn-Tucker (KKT)). *Let the cost function f and constraint functions g_i for all $i \in [m]$ be convex and differentiable functions, such that the constraints are qualified. Then $x^* \in \mathbb{R}^N$ is a solution of the constrained problem iff there exists $\alpha^* \in \mathbb{R}_+^m$ such that*

$$\nabla_x \mathcal{L}(x^*, \alpha^*) = \nabla_x f(x^*) + \langle \alpha^*, \nabla_x g(x^*) \rangle = 0, \quad \nabla_\alpha \mathcal{L}(x^*, \alpha^*) = g(x^*) \leq 0, \quad \langle \alpha^*, g(x^*) \rangle = 0.$$

Proof. From the necessary condition theorem, it follows that if x^* is a solution to the primal problem, then there exists dual variables α^* such that (x^*, α^*) is a saddle point of the Lagrangian, and all three conditions are satisfied.

Conversely, if all three conditions are met, then for any $x \in \mathbb{R}^N$ such that $g_i(x) \leq 0$ for all $i \in [m]$, we have

$$f(x) - f(x^*) \geq \langle \nabla_x f(x^*), x - x^* \rangle = - \sum_{i=1}^m \alpha_i^* \langle \nabla_x g_i(x^*), x - x^* \rangle \geq - \langle \alpha^*, g(x) - g(x^*) \rangle = - \langle \alpha^*, g(x) \rangle \geq 0.$$

The first inequality follows from the convexity of f . The subsequent equality follows from the first condition. Next inequality follows from the convexity of g_i for all $i \in [m]$. Next equality follows from the third condition, and the last inequality from the fact that $x \in \mathcal{X}$. \square