

# Lecture-08: Expectation

## 1 Expectation

**Example 1.1.** Consider a probability space  $(\Omega, \mathcal{F}, P)$ . We consider  $N$  trials of a random experiment, and define a random vector  $X : \Omega \rightarrow \mathcal{X}^N$  such that  $X_i(\omega)$  is a discrete random variable associated with the trial  $i \in [N]$ . If the marginal distributions of random variables  $(X_i : \Omega \rightarrow \mathcal{X} : i \in [N])$  are identical with the common probability mass function  $P_{X_1} : \mathcal{X} \rightarrow [0, 1]$ , then the empirical mean of random variable  $X_1$  can be written as

$$\hat{m} = \frac{1}{N} \sum_{i=1}^N X_i(\omega).$$

For a random variable  $X_1 : \Omega \rightarrow \mathcal{X}$ , we can define events  $C_{X_1}(x) \triangleq X_1^{-1}\{x\}$  for each value  $x \in \mathcal{X}$ . The probability mass function  $P_{X_1} : \mathcal{X} \rightarrow [0, 1]$  for a discrete random variable can be estimated for each  $x \in \mathcal{X}$  as the empirical probability mass function

$$\hat{P}_{X_1}(x) = \frac{1}{N} \sum_{i=1}^N \mathbb{1}_{C_{X_1}(x)}(\omega).$$

Recall that a simple random variable  $X_1$  can be written as  $X_1 = \sum_{x \in \mathcal{X}} x \mathbb{1}_{C_{X_1}(x)}$ , where  $C_{X_1} \triangleq (C_{X_1}(x) \in \mathcal{F} : x \in \mathcal{X})$  is a finite partition of the sample space  $\Omega$  and  $P_{X_1}(x) = P(C_{X_1}(x))$ . That is, we can write the empirical mean in terms of the empirical PMF as

$$\hat{m} = \frac{1}{N} \sum_{i=1}^N \sum_{x \in \mathcal{X}} x \mathbb{1}_{C_{X_1}(x)}(\omega) = \sum_{x \in \mathcal{X}} x \hat{P}_{X_1}(x).$$

This example motivates the following definition of mean for simple random variables.

**Definition 1.2 (Expectation of simple random variable).** The **mean** or **expectation** of a simple random variable  $X : \Omega \rightarrow \mathcal{X} \subseteq \mathbb{R}$  defined on a probability space  $(\Omega, \mathcal{F}, P)$ , is denoted by  $\mathbb{E}[X]$  and defined as

$$\mathbb{E}[X] \triangleq \sum_{x \in \mathcal{X}} x P_X(x).$$

*Remark 1.* Since a simple random variable can be written as  $X = \sum_{x \in \mathcal{X}} x \mathbb{1}_{\{X=x\}}$  for  $C_X(x) = X^{-1}\{x\}$ , we can write the expectation of a simple random variable as an integral

$$\mathbb{E}[X] = \int_{\Omega} X(\omega) P(d\omega) = \int_{\Omega} \sum_{x \in \mathcal{X}} x \mathbb{1}_{C_X(x)}(\omega) P(d\omega) = \sum_{x \in \mathcal{X}} x \int_{\Omega} \mathbb{1}_{C_X(x)}(\omega) P(d\omega) = \sum_{x \in \mathcal{X}} x \mathbb{E}[\mathbb{1}_{C_X(x)}] = \sum_{x \in \mathcal{X}} x P_X(x).$$

That is, the expectation of an indicator function is the probability of the indicated set.

**Theorem 1.3.** Consider a non-negative random variable  $X : \Omega \rightarrow \mathbb{R}_+$  defined on a probability space  $(\Omega, \mathcal{F}, P)$ . There exists a sequence of non-decreasing non-negative simple random variables  $Y : \Omega \rightarrow \mathbb{R}_+^{\mathbb{N}}$  such that for all  $\omega \in \Omega$

$$Y_n(\omega) \leq Y_{n+1}(\omega), \text{ for all } n \in \mathbb{N}, \text{ and } \lim_n Y_n(\omega) = X(\omega).$$

Then  $\mathbb{E}[Y_n]$  is defined for each  $n \in \mathbb{N}$ , the sequence  $(\mathbb{E}[Y_n] \in \mathbb{R}_+ : n \in \mathbb{N})$  is non-decreasing, and the limit  $\lim_n \mathbb{E}[Y_n] \in \mathbb{R}_+ \cup \{\infty\}$  exists. This limit is independent of the choice of the sequence and depends only on the probability space.

*Proof.* For each  $n \in \mathbb{N}$  and  $k \in \{0, \dots, 2^n - 1\}$ , we define half-open sets  $B_{n,k} \triangleq (k2^{-n}, (k+1)2^{-n}]$ . Then, the collection of sets  $B_n \triangleq (B_{n,k} : k \in \{0, \dots, 2^n - 1\})$  partitions the set  $(0, 2^n]$  for each  $n \in \mathbb{N}$ . Further, we observe that  $\cup_{n \in \mathbb{N}} (0, 2^n] = \mathbb{R}^+$  and that  $B_{n+1,2k} \cup B_{n+1,2k+1} = B_{n,k}$  for all  $n \in \mathbb{N}$  and  $k$ .

For a non-negative random variable  $X : \Omega \rightarrow \mathbb{R}_+$ , we define events  $A_{n,k}^X = X^{-1}(B_{n,k}) \in \mathcal{F}$ , and a sequence of simple non-negative random variables  $Y : \Omega \rightarrow \mathbb{R}_+^{\mathbb{N}}$  in the following fashion

$$Y_n(\omega) \triangleq \sum_{k=0}^{2^n-1} \mathbb{1}_{A_{n,k}^X}(\omega) \left( \inf_{\omega \in A_{n,k}^X} X(\omega) \right) = \sum_{k=0}^{2^n-1} \mathbb{1}_{A_{n,k}^X}(\omega) \left( \inf_{X(\omega) \in B_{n,k}} X(\omega) \right) = \sum_{k=0}^{2^n-1} k2^{-n} \mathbb{1}_{A_{n,k}^X}(\omega).$$

We observe that  $Y_n$  is a quantized version of  $X$ , and its value is the left end-point  $k2^{-n}$  when  $X \in B_{n,k}$  for each  $k \in \{0, \dots, 2^n - 1\}$ . Since  $\cup_{k=0}^{2^n-1} A_{n,k}^X = X^{-1}(0, 2^n]$ , it follows that we cover the positive real line as  $n$  grows larger and the step size grows smaller. Thus, the limiting random variable can take all possible non-negative real values. We observe that

$$\begin{aligned} Y_{n+1}(\omega) &= \sum_{k=0}^{2^{2(n+1)}-1} \mathbb{1}_{A_{n+1,k}^X}(\omega) \left( \inf_{X(\omega) \in B_{n+1,k}} X(\omega) \right) \\ &= \sum_{k=0}^{2^{2n}-1} \left( \mathbb{1}_{A_{n+1,2k}^X}(\omega) \left( \inf_{X(\omega) \in B_{n+1,2k}} X(\omega) \right) + \mathbb{1}_{A_{n+1,2k+1}^X}(\omega) \left( \inf_{X(\omega) \in B_{n+1,2k+1}} X(\omega) \right) \right) \\ &\geq \sum_{k=0}^{2^{2n}-1} \mathbb{1}_{A_{n,2k}^X}(\omega) \left( \inf_{X(\omega) \in B_{n,k}} X(\omega) \right) = Y_n(\omega). \end{aligned}$$

We see that  $Y_n(\omega) \leq Y_{n+1}(\omega) \leq X(\omega)$  and  $\lim_n Y_n(\omega) = X(\omega)$  for all  $\omega \in \Omega$ .

Since  $Y_n : \Omega \rightarrow \mathbb{R}$  is a simple random variable for all  $n \in \mathbb{N}$ , the expectation  $\mathbb{E}[Y_n]$  is defined for all  $n$ , and can be written as

$$\mathbb{E}[Y_n] = \sum_{k=0}^{2^n-1} k2^{-n} [F_X((k+1)2^{-n}) - F_X(k2^{-n})].$$

We observe that this expectation is completely specified by the distribution function  $F_X$ , and we can write the limit

$$\lim_n \mathbb{E}[Y_n] = \lim_n \sum_{k=0}^{2^n-1} k2^{-n} [F_X(k2^{-n} + 2^{-n}) - F_X(k2^{-n})] = \int_{\mathbb{R}^+} x dF_X(x).$$

□

**Definition 1.4 (Expectation of a non-negative random variable).** For a non-negative random variable  $X : \Omega \rightarrow \mathbb{R}$  defined on the probability space  $(\Omega, \mathcal{F}, P)$ , consider the sequence of non-decreasing simple random variables  $Y : \Omega \rightarrow \mathbb{R}_+^{\mathbb{N}}$  such that  $\lim_n Y_n = X$ . The **expectation** of the non-negative random variable  $X$  is defined as

$$\mathbb{E}[X] \triangleq \lim_n \mathbb{E}[Y_n].$$

*Remark 2.* From the definition, it follows that  $\mathbb{E}[X] = \int_{\mathbb{R}^+} x dF_X(x)$ .

**Definition 1.5 (Expectation of a real random variable).** For a real-valued random variable  $X$  defined on a probability space  $(\Omega, \mathcal{F}, P)$ , we can define the following functions

$$X_+ \triangleq \max\{X, 0\}, \quad X_- \triangleq \max\{0, -X\}.$$

We can verify that  $X_+, X_-$  are non-negative random variables and hence their expectations are well defined. We observe that  $X(\omega) = X_+(\omega) - X_-(\omega)$  for each  $\omega \in \Omega$ . If at least one of the  $\mathbb{E}[X_+]$  and  $\mathbb{E}[X_-]$  is finite, then the **expectation** of the random variable  $X$  is defined as

$$\mathbb{E}[X] \triangleq \mathbb{E}[X_+] - \mathbb{E}[X_-].$$

**Theorem 1.6 (Expectation as an integral with respect to the distribution function).** For a random variable  $X : \Omega \rightarrow \mathbb{R}$  defined on the probability space  $(\Omega, \mathcal{F}, P)$ , the expectation is given by

$$\mathbb{E}[X] = \int_{\mathbb{R}} x dF_X(x).$$

*Proof.* It suffices to show this for a non-negative random variable  $X$ , and the result follows from the definition of expectation of a non-negative random variable as the limit of expectation of approximating simple functions.  $\square$

## 2 Properties of Expectations

**Theorem 2.1 (Properties).** Let  $X : \Omega \rightarrow \mathbb{R}$  be a random variable defined on the probability space  $(\Omega, \mathcal{F}, P)$ .

(i) **Linearity:** Let  $a, b \in \mathbb{R}$  and  $X, Y$  be random variables defined on the probability space  $(\Omega, \mathcal{F}, P)$ . If  $\mathbb{E}X, \mathbb{E}Y$ , and  $a\mathbb{E}X + b\mathbb{E}Y$  are well defined, then  $\mathbb{E}(aX + bY)$  is well defined and

$$\mathbb{E}(aX + bY) = a\mathbb{E}X + b\mathbb{E}Y.$$

(ii) **Monotonicity:** If  $P\{X \geq Y\} = 1$  and  $\mathbb{E}[Y]$  is well defined with  $\mathbb{E}[Y] > -\infty$ , then  $\mathbb{E}[X]$  is well defined and  $\mathbb{E}[X] \geq \mathbb{E}[Y]$ .

(iii) **Functions of random variables:** Let  $g : \mathbb{R} \rightarrow \mathbb{R}$  be a Borel measurable function, then  $g(X)$  is a random variable with  $\mathbb{E}[g(X)] = \int_{x \in \mathbb{R}} g(x) dF(x)$ .

(iv) **Continuous random variables:** Let  $f_X : \mathbb{R} \rightarrow [0, \infty)$  be the density function, then  $\mathbb{E}X = \int_{x \in \mathbb{R}} x f_X(x) dx$ .

(v) **Discrete random variables:** Let  $P_X : \mathcal{X} \rightarrow [0, 1]$  be the probability mass function, then  $\mathbb{E}X = \sum_{x \in \mathcal{X}} x P_X(x)$ .

(vi) **Integration by parts:** The expectation  $\mathbb{E}X = \int_{x \geq 0} (1 - F_X(x)) dx + \int_{x < 0} F_X(x) dx$  is well defined when at least one of the two parts is finite on the right hand side.

*Proof.* It suffices to show properties (i) – (iii) for simple random variables.

(i) Let  $X = \sum_{x \in \mathcal{X}} x \mathbb{1}_{C_X(x)}$  and  $Y = \sum_{y \in \mathcal{Y}} y \mathbb{1}_{C_Y(y)}$  be simple random variables, then  $(C_X(x) \cap C_Y(y) \in \mathcal{F} : (x, y) \in \mathcal{X} \times \mathcal{Y})$  partition the sample space  $\Omega$ . Hence, we can write  $aX + bY = \sum_{(x, y) \in \mathcal{X} \times \mathcal{Y}} (ax + by) \mathbb{1}_{\{C_X(x) \cap C_Y(y)\}}$  and from linearity of sum it follows that

$$\begin{aligned} \mathbb{E}[aX + bY] &= \sum_{(x, y) \in \mathcal{X} \times \mathcal{Y}} (ax + by) P\{C_X(x) \cap C_Y(y)\} \\ &= a \sum_{x \in \mathcal{X}} x \sum_{y \in \mathcal{Y}} P\{C_X(x) \cap C_Y(y)\} + b \sum_{y \in \mathcal{Y}} y \sum_{x \in \mathcal{X}} P\{C_X(x) \cap C_Y(y)\} \\ &= a \sum_{x \in \mathcal{X}} x P(C_X(x)) + b \sum_{y \in \mathcal{Y}} y P(C_Y(y)) = a\mathbb{E}X + b\mathbb{E}Y. \end{aligned}$$

(ii) From the fact that  $X - Y \geq 0$  almost surely and linearity of expectation, it suffices to show that  $\mathbb{E}X \geq 0$  for non-negative random variable  $X$ . It can easily be shown for simple non-negative random variables, and follows for general non-negative random variables by taking limits.

(iii) It suffices to show this holds true for simple random variables  $X : \Omega \rightarrow \mathcal{X} \subset \mathbb{R}$ . Since  $g : \mathbb{R} \rightarrow \mathbb{R}$  is Borel measurable,  $Y = g(X)$  is a random variable. For each  $y \in \mathcal{Y} = g(\mathcal{X})$ , we have

$$C_Y(y) = \{\omega \in \Omega : (g \circ X)(\omega) = y\} = X^{-1} \circ g^{-1}\{y\} = \cup_{x \in g^{-1}\{y\}} C_X(x).$$

Therefore, we can write the expectation

$$\mathbb{E}[Y] = \sum_{y \in \mathcal{Y}} y P(C_Y(y)) = \sum_{y \in \mathcal{Y}} \sum_{x \in g^{-1}(y)} g(x) P(C_X(x)) = \sum_{x \in \mathcal{X}} g(x) P(C_X(x)).$$

(iv) For continuous random variables, we have  $dF_X(x) = f_X(x)dx$  for all  $x \in \mathbb{R}$ .

(v) For discrete random variables  $X : \Omega \rightarrow \mathcal{X}$ , we have  $dF_X(x) = P_X(x)$  for all  $x \in \mathcal{X}$  and zero otherwise.

(vi) We can write  $\mathbb{E}X = -\int_{x \geq 0} xd(1 - F_X)(x) + \int_{x < 0} xdF_X(x)$ . Therefore, we have

$$= -x(1 - F_X(x))\Big|_0^\infty + \int_{x \geq 0} (1 - F_X(x))dx.$$

□