

Lecture-22: Communicating classes

1 Communicating classes

Definition 1.1. For states $x, y \in \mathcal{X}$, it is said that state y is **accessible** from state x if $p_{xy}^{(n)} > 0$ for some $n \in \mathbb{Z}_+$, and denoted by $x \rightarrow y$. If two states $x, y \in \mathcal{X}$ are accessible to each other, they are said to **communicate** with each other, denoted by $x \leftrightarrow y$.

Proposition 1.2. *Communication is an equivalence relation.*

Proof. Relation on state space \mathcal{X} is a subset of product of sets $\mathcal{X} \times \mathcal{X}$. Communication is a relation on state space \mathcal{X} , as it relates two states $x, y \in \mathcal{X}$. To show equivalence, we have to show reflexivity, symmetry, and transitivity of the relation.

Reflexivity: Since $P^0 = I$, it follows that $p_{xx}^{(0)} = 1 > 0$ for each state $x \in \mathcal{X}$, and hence $x \leftrightarrow x$.

Symmetry: If $x \leftrightarrow y$, then we know that $x \rightarrow y$ and $y \rightarrow x$ and hence $y \leftrightarrow x$.

Transitivity: For transitivity, suppose $x \leftrightarrow y$ and $y \leftrightarrow z$. Let $m, n \in \mathbb{Z}_+$ such that $p_{xy}^{(m)} > 0$ and $p_{yz}^{(n)} > 0$. Then by Chapman Kolmogorov equation, we have $p_{xz}^{(m+n)} = \sum_{w \in \mathcal{X}} p_{xw}^{(m)} p_{wz}^{(n)} \geq p_{xy}^{(m)} p_{yz}^{(n)} > 0$. This implies $x \rightarrow z$, and using similar arguments one can show that $z \rightarrow x$, and the transitivity follows. □

Remark 1. Since the communication is an equivalence relation, it partitions state space \mathcal{X} into equivalence classes.

Definition 1.3. Each equivalence class of communication relation is called a **communicating class**. A property of states is said to be a **class property** if for each communicating class \mathcal{C} , either all states in \mathcal{C} have the property, or none do.

Definition 1.4. A set of states that communicate are called a **communicating class**.

- (i) A communicating class \mathcal{C} is called **closed** if no edges leave this class. That is, for all edges $(x, y) \in \mathcal{C} \times \mathcal{C}^c$, we have $p_{xy} = 0$.
- (ii) An **open** communicating class is not closed, i.e. there exist an edge that leaves this class. That is, there exists an edge $(x, y) \in \mathcal{C} \times \mathcal{C}^c$ such that $p_{xy} > 0$.

1.1 Irreducibility and periodicity

Definition 1.5. A Markov chain with a single class is called an **irreducible** Markov chain. That is, for any two states $x, y \in \mathcal{X}$, there exists an integer $n \in \mathbb{Z}_+$ such that $p_{xy}^{(n)} > 0$. In other words, any state y can be reached from any state x using transitions of positive probability.

Definition 1.6. Let $\mathcal{T}(x) \triangleq \{n \in \mathbb{N} : p_{xx}^{(n)} > 0\}$ be the set of times when the chain can possibly return to the initial state x . The **period** of any state $x \in \mathcal{X}$ is defined as $d(x) \triangleq \gcd \mathcal{T}(x) = \gcd\{n \in \mathbb{N} : p_{xx}^{(n)} > 0\}$. We define $d(x) = \infty$, if $p_{xx}^{(n)} = 0$ for all $n \in \mathbb{N}$. A state $x \in \mathcal{X}$ is called **aperiodic** if the period $d(x)$ is 1.

Proposition 1.7. *If $x \leftrightarrow y$, then $d(x) = d(y)$. That is, periodicity is a class property.*

Proof. Consider two states $x \neq y \in \mathcal{X}$ and integers $m, n \in \mathbb{N}$ be such that $p_{xy}^{(m)} p_{yx}^{(n)} > 0$. Suppose $s \in \mathcal{T}(x)$, that is $p_{xx}^{(s)} > 0$. Then

$$p_{yy}^{(n+m)} \geq p_{yx}^{(n)} p_{xy}^{(m)} > 0, \quad p_{yy}^{(n+s+m)} \geq p_{yx}^{(n)} p_{xx}^{(s)} p_{xy}^{(m)} > 0.$$

Hence $d(y)|n+m$ and $d(y)|n+s+m$, and hence $d(y)|s$ for any $s \in \mathcal{T}(x)$. In particular, it implies that $d(y)|d(x)$. By symmetrical arguments, we get $d(x)|d(y)$. Hence $d(x) = d(y)$. \square

Definition 1.8. For an irreducible chain, the period of the chain is defined to be the period which is common to all states. An irreducible Markov chain is called **aperiodic** if the single communicating class is aperiodic.

Proposition 1.9. *If the transition matrix P is aperiodic and irreducible, then there is an integer r_0 such that $p_{xy}^{(r)} > 0$ for all $x, y \in \mathcal{X}$ and $r \geq r_0$.*

1.2 Transient and recurrent states

Proposition 1.10. *Transience and recurrence are class properties.*

Proof. Let us start with proving recurrence is a class property. Let x be a recurrent state and let $x \leftrightarrow y$. Hence there exist some $m, n > 0$, such that $p_{xy}^{(m)} > 0$ and $p_{yx}^{(n)} > 0$. As a consequence of the recurrence, $\sum_{s \in \mathbb{N}} p_{xx}^{(s)} = \infty$. $\sum_{s \in \mathbb{N}} p_{yy}^{(m+n+s)} \geq \sum_{s \in \mathbb{N}} p_{yx}^{(n)} p_{xx}^{(s)} p_{xy}^{(m)} = \infty$. If x were transient instead then $\sum_{s \in \mathbb{N}} p_{xx}^{(s)} < \infty$, and we conclude that y is also transient by the following observation $\sum_{s \in \mathbb{N}} p_{yy}^{(s)} \leq \frac{\sum_{s \in \mathbb{N}} p_{xx}^{(m+n+s)}}{p_{yx}^{(n)} p_{xy}^{(m)}} < \infty$. \square

Corollary 1.11. *If y is a recurrent state and there exists a state x such that $y \rightarrow x$, then $x \rightarrow y$ and $f_{xy} = 1$.*

Proof. Let $y \in \mathcal{X}$ be a recurrent state, and consider state $x \in \mathcal{X}$ such that $y \rightarrow x$. We will show that $f_{xy} = 1$ and hence $f_{xy}^{(n)} > 0$ for some $n \in \mathbb{Z}_+$ and $x \rightarrow y$. To this end, we observe that since $y \rightarrow x$, there exists an integer $n \in \mathbb{Z}_+$ such that the probability of hitting state x for the first time starting from state y in n -steps is positive. That is,

$$f_{yx}^{(n)} \triangleq P_y \{X_n = x, X_{n-1} \neq x, \dots, X_1 \neq x\} = P_y \left\{ \tau_X^{\{x\},1} = n \right\} > 0.$$

From the strong Markov property, we have

$$1 - f_{yy} = P_y \left\{ \tau_X^{\{y\},1} = \infty \right\} \geq P_y \left\{ \tau_X^{\{y\},1} = \infty, \tau_X^{\{x\},1} = n \right\} = P_x \left\{ \tau_X^{\{y\},1} = \infty \right\} P_y \left\{ \tau_X^{\{x\},1} = n \right\} = f_{yx}^{(n)} (1 - f_{xy}).$$

Since state y is recurrent, it follows that $f_{xy} = 1$ and hence $x \rightarrow y$. \square

Corollary 1.12. *Let $x, y \in \mathcal{X}$ be in the same communicating class and the state y is recurrent. Then, $\lim_{n \in \mathbb{N}} \frac{\sum_{k=1}^n p_{xy}^{(k)}}{n} = \frac{1}{\mu_{yy}}$. Furthermore, if the state y is aperiodic, then $\lim_{n \in \mathbb{N}} p_{xy}^{(n)} = \frac{1}{\mu_{yy}}$.*

Proof. Since y is recurrent and $y \rightarrow x$, it follows that $f_{xy} = 1$ from the previous Lemma. From the Theorem 1.7 in previous lecture, it follows that $\lim_{n \in \mathbb{N}} \frac{\sum_{k=1}^n p_{xy}^{(k)}}{n} = \frac{1}{\mu_{yy}}$.

Let the period of the state y be d . Then we know that there exists a positive integer r_0 such that for all $n \geq r_0$, we have $p_{yy}^{(nd)} > 0$. \square

Theorem 1.13. *The states in a communicating class are of one of the following types; all transient, or all null recurrent, or all positive recurrent.*

Proof. It suffices to show that if x, y belong to the same communicating class and y is null recurrent, then x is null recurrent as well. We take $r, s \in \mathbb{N}$, such that $p_{yx}^{(r)} p_{xy}^{(s)} > 0$. It follows that $p_{yy}^{(r+s)} \geq p_{yx}^{(r)} p_{xx}^{(s)} p_{xy}^{(s)}$ for all $\ell \in \mathbb{N}$. Hence, for any $n > r + s$, we have

$$\frac{1}{n} \sum_{k=1}^n p_{yy}^{(k)} \geq \frac{1}{n} \sum_{k=r+s+1}^n p_{yy}^{(k)} \geq \left(\frac{n-r-s}{n} \right) \left(\frac{1}{n-r-s} \sum_{\ell=1}^{n-r-s} p_{xx}^{(\ell)} \right) p_{yx}^{(r)} p_{xy}^{(s)}.$$

Since y is null recurrent LHS goes to zero as n increases, which implies $\lim_{n \in \mathbb{N}} \frac{1}{n} \sum_{\ell=1}^n p_{xx}^{(\ell)} = 0$. Hence, x is null recurrent as well. \square