

# Lecture-11: Key Lemma and Blackwell Theorem

## 1 Key Lemma

**Theorem 1.1 (Key Lemma).** Consider a renewal sequence  $S : \Omega \rightarrow \mathbb{R}_+^{\mathbb{N}}$  with i.i.d. inter-renewal times  $X : \Omega \rightarrow \mathbb{R}_+^{\mathbb{N}}$  having common distribution function  $F$ , associated counting process  $N : \Omega \rightarrow \mathbb{Z}_+^{\mathbb{R}_+}$ , and the renewal function  $m : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ . Then,

$$P\{S_{N_t} \leq s\} = \bar{F}(t) + \int_0^s \bar{F}(t-y) dm(y), \quad t \geq s \geq 0.$$

*Proof.* We can see that event of time of last renewal prior to  $t$  being smaller than another time  $s$  can be partitioned into disjoint events corresponding to number of renewals until time  $t$ . Each of these disjoint events is equivalent to occurrence of  $n$ th renewal before time  $s$  and  $(n+1)$ th renewal past time  $t$ . That is,

$$\{S_{N_t} \leq s\} = \bigcup_{n \in \mathbb{Z}_+} \{S_{N_t} \leq s, N_t = n\} = \bigcup_{n \in \mathbb{Z}_+} \{S_n \leq s, S_{n+1} > t\}.$$

Recognizing that  $S_0 = 0, S_1 = X_1$ , and that  $S_{n+1} = S_n + X_{n+1}$ , we can write

$$P\{S_{N_t} \leq s\} = P\{X_1 > t\} + \sum_{n \in \mathbb{N}} \mathbb{E}[\mathbb{1}_{\{S_n \leq s\}} \mathbb{E}[\mathbb{1}_{\{X_{n+1} > t - S_n\}} | \sigma(S_n)]].$$

We recall that the distribution function of  $n$ th renewal instant  $S_n$  is the  $n$ -fold convolution of  $F$  denoted by  $F_n$ . Taking expectation of  $\bar{F}(t - S_n) \mathbb{1}_{\{S_n \leq s\}}$ , we get

$$P\{S_{N_t} \leq s\} = \bar{F}(t) + \sum_{n \in \mathbb{N}} \int_{y=0}^s \bar{F}(t-y) dF_n(y).$$

Using monotone convergence theorem to interchange integral and summation, and noticing that  $m(y) = \sum_{n \in \mathbb{N}} F_n(y)$ , the result follows.  $\square$

*Remark 1.* Key lemma tells us that distribution of  $S_{N_t}$  has probability mass at 0 and density between  $(0, t]$ , that is,

$$P\{S_{N_t} = 0\} = \bar{F}(t), \quad dF_{S_{N_t}}(y) = \bar{F}(t-y) dm(y), \quad 0 < y \leq t.$$

*Remark 2.* Density of  $S_{N_t}$  has interpretation of renewal taking place in the infinitesimal neighborhood of  $y$ , and next inter-arrival after time  $t - y$ . To see this, we notice

$$dm(y) = \sum_{n \in \mathbb{N}} dF_n(y) = \sum_{n \in \mathbb{N}} P\{S_n \in (y, y + dy)\} = \sum_{n \in \mathbb{N}} P\{n\text{th renewal occurs in } (y, y + dy)\}.$$

Combining interpretation of density of inter-arrival time  $dF(t)$ , we get

$$dF_{S_{N_t}}(y) = P\{\text{renewal occurs in } (y, y + dy) \text{ and next arrival after } t - y\}. \quad (1)$$

**Example 1.2 (Poisson process).** Let the inter-renewal time i.i.d. random sequence  $X : \Omega : \mathbb{R}_+^{\mathbb{N}}$  be exponentially distributed with common distribution  $F(x) = 1 - e^{-\lambda x}$  for  $x \in \mathbb{R}_+$ . Then, the distribution of last renewal is given by

$$P\{S_{N_t} \leq x\} = e^{-\lambda t} + \int_0^x \lambda e^{-\lambda(t-y)} dy = e^{-\lambda(t-x)}, \quad 0 \leq x \leq t.$$

## 2 Delayed Regenerative Process

**Theorem 2.1.** Let  $Z$  be a delayed regenerative process with the associated delayed renewal sequence  $S$ , the renewal function  $m_D$ , the first arrival distribution  $G$ , and the common inter-arrival duration distribution  $F$ . For a Borel measurable set  $A \in \mathcal{B}(\mathbb{R})$ , we define the kernel functions  $K_1(t) \triangleq P\{Z_t \in A, S_1 > t\}$ ,  $K_2(t) \triangleq P\{Z_{S_1+t} \in A, t \in [0, X_2)\}$ , then we have

$$P\{Z_t \in A\} = K_1(t) + \int_0^t dm_D(y)K_2(t-y). \quad (2)$$

*Proof.* For a Borel measurable set  $A \in \mathcal{B}(\mathbb{R})$ , we can write the probability of the delayed regenerative process taking values in this set as disjoint sum of probability of disjoint partitions of this event as

$$P\{Z_t \in A\} = P\{Z_t \in A, S_1 > t\} + \sum_{n \in \mathbb{N}} P\{Z_t \in A, N_t = n\}.$$

The  $n$ th segment of the joint process  $(N_t^D, Z_t)$  is  $\zeta_n = (X_n, (Z_{S_{n-1}+t} : t \in [0, X_n)))$ . From the regenerative property, we know that the segments  $(\zeta_n : n \in \mathbb{N})$  are independent, where  $(\zeta_n : n \geq 2)$  are identically distributed. In particular, we can write

$$\mathbb{E}[\mathbb{1}_{\{Z_t \in A, S_n \leq t < S_{n+1}\}} | \mathcal{F}_{S_n}] = \mathbb{1}_{\{S_n \leq t\}} \mathbb{E}[\mathbb{1}_{\{Z_{t-S_n} \in A, t-S_n \in [0, X_{n+1})\}} | \sigma(S_n)] = \mathbb{1}_{\{S_n \leq t\}} K_2(t - S_n).$$

The result follows from the fact that  $P\{Z_t \in A, N_t = n\} = \mathbb{E}[\mathbb{1}_{\{Z_t \in A, S_n \leq t < S_{n+1}\}}]$ .  $\square$

**Example 2.2 (Age process).** Age process  $(A(t) = t - S_{N(t)} : t \geq 0)$  for a delayed renewal process  $(S_n : n \in \mathbb{N})$  is a delayed regenerative process, since the  $n$ th segment is given by  $\zeta_n = (X_n, (A(S_{n-1} + t) = t : t \geq [0, X_n)))$ . For the measurable set  $B = [x, \infty)$ , then we can compute the kernel functions

$$K_1(t) = P\{A(t) \geq x, S_1 > t\} = \mathbb{1}_{\{t \geq x\}} \bar{G}(t), \quad K_2(t) = P\{A(S_1 + t) \geq x, t \in [0, X_2)\} = \mathbb{1}_{\{t \geq x\}} \bar{F}(t).$$

Therefore, we can write the distribution of last renewal time for the delayed renewal process as

$$P\{S_{N(t)} \leq x\} = P\{A(t) \geq t - x\} = \mathbb{1}_{\{x \geq 0\}} \bar{G}(t) + \int_0^t dm_D(y) \mathbb{1}_{\{t-y \geq t-x\}} \bar{F}(t-y).$$

**Corollary 2.3 (Delayed Key Lemma).** Consider a delayed renewal sequence  $S : \Omega \rightarrow \mathbb{R}_+^{\mathbb{N}}$  with independent inter-renewal times  $X : \Omega \rightarrow \mathbb{R}_+^{\mathbb{N}}$  with first renewal time having distribution  $G$  and common distribution  $F$  for inter-renewal times  $(X_n, n \geq 2)$ , associated counting process  $N^D : \Omega \rightarrow \mathbb{Z}_+^{\mathbb{R}_+}$ , and the renewal function  $m_D : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ . Then,

$$P\{S_{N_t^D} \leq s\} = \bar{G}(t) + \int_0^s \bar{F}(t-y) dm_D(y), \quad t \geq s \geq 0.$$

## 3 Blackwell Theorem

**Lemma 3.1.** For a renewal sequence  $S$ , let  $F$  be the inter-renewal time distribution such that  $\inf\{x : F(x) = 1\} = \infty$ , then for any  $b > 0$

$$\sup_t \{m_t - m_{t-b}\} < \infty.$$

*Proof.* Recall that  $m = \sum_{n \in \mathbb{N}} F_n$  and hence  $m * F = m - F$ . This implies that  $m * (1 - F) = F$ . Since the function  $1 - F$  is monotonically non-increasing,  $\inf_{s \in [0, b]} \bar{F}(s) = \bar{F}(b)$ . Therefore,

$$1 \geq F(t) = \int_0^t dm(s) \bar{F}(t-s) \geq \int_{t-b}^t dm(s) \bar{F}(t-s) \geq [m_t - m_{t-b}] \bar{F}(b),$$

where  $b$  is chosen so that  $F(b) < 1$ . Hence, the result follows.  $\square$

**Theorem 3.2 (Blackwell's Theorem).** Consider a renewal sequence  $S$  with the inter-renewal time distribution  $F$  such that  $\inf\{x : F(x) = 1\} = \infty$ , mean of inter-renewal time  $\mu$ , and renewal function  $m_t$ . If  $F$  is not lattice, then for all  $a \geq 0$

$$\lim_{t \rightarrow \infty} m_{t+a} - m_t = \frac{a}{\mu}.$$

If  $F$  is lattice with period  $d$ , then

$$\lim_{n \rightarrow \infty} m_{(n+1)d} - m_{nd} = \frac{d}{\mu}.$$

*Proof.* We will not prove that the following limit exists for non-lattice  $F$ ,

$$g(a) \triangleq \lim_{t \rightarrow \infty} [m_{t+a} - m_t] \quad (3)$$

However, we show that if this limit does exist, it is equal to  $a/\mu$  as a consequence of elementary renewal theorem. To this end, note that

$$m_{t+a+b} - m_t = m_{t+a+b} - m_{t+a} + m_{t+a} - m_t.$$

Taking limits on both sides of the above equation, we conclude that  $g(a+b) = g(a) + g(b)$ . The only increasing solution of such a  $g : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  for all  $a > 0$  is

$$g(a) = ca,$$

for some positive constant  $c$ . To show  $c = \frac{1}{\mu}$ , define a sequence  $x \in \mathbb{R}_+^{\mathbb{N}}$  in terms of renewal function  $m_t$ , as

$$x_n \triangleq m_n - m_{n-1}, \quad n \in \mathbb{N}.$$

Note that  $\sum_{i=1}^n x_i = m_n$  and  $\lim_{n \in \mathbb{N}} x_n = g(1) = c$ , hence we have the Cesàro mean converging to

$$\lim_{n \in \mathbb{N}} \frac{\sum_{i=1}^n x_i}{n} = \lim_{n \in \mathbb{N}} \frac{m_n}{n} \stackrel{(a)}{=} c,$$

where (a) follows from the fact that if a sequence  $\{x_i\}$  converges to  $c$ , then the running average sequence  $a_n = \frac{1}{n} \sum_{i=1}^n x_i$  also converges to  $c$ , as  $n \rightarrow \infty$ . Therefore, we can conclude  $c = 1/\mu$  by elementary renewal theorem.

When  $F$  is lattice with period  $d$ , the limit in (3) doesn't exist. (See the following example). However, the theorem is true for lattice again by elementary renewal theorem. Indeed, since  $\frac{m_{nd}}{n} \rightarrow \frac{1}{\mu}$ , we can define  $x_n \triangleq m_{nd} - m_{(n-1)d}$  and observe that  $\sum_{i=1}^n x_i = m_{nd}$  and the Cesàro mean  $\frac{1}{n} \sum_{i=1}^n x_i$  converges to  $\frac{d}{\mu}$  by elementary renewal theorem.  $\square$

**Example 3.3.** Consider a renewal process with  $P\{X_n = 1\} = 1$ , that is, there is a renewal at every positive integer time instant with unit probability. Then  $F$  is lattice with  $d = 1$ . Now, for  $a = 0.5$ , and  $t_n = n + (-1)^n 0.5$ , we see that  $m_{t_n} = N_{t_n} = n - \mathbb{1}_{\{n \text{ odd}\}}$ , and  $m(t_n + a) = n$ . It follows that  $m_{t_n+a_n} - m_{t_n} = \mathbb{1}_{\{n \text{ odd}\}}$ , and hence  $\lim_{t_n \rightarrow \infty} m_{t_n+a} - m_{t_n}$  does not exist. It follows that  $\lim_{t \rightarrow \infty} m_{t+a} - m_t$  does not exist.

*Remark 3.* In the lattice case, if the inter arrivals are strictly positive, that is, there can be no more than one renewal at each  $nd$ , then we have that

$$\lim_{n \rightarrow \infty} P\{\text{renewal at } nd\} = \frac{d}{\mu}.$$

**Corollary 3.4 (Delayed Blackwell's Theorem).** Consider a delayed renewal process with independent inter-renewal times, with the distribution of first renewal being  $G$  with mean  $\mu_G$ , and the distribution of inter-renewal times for  $n \geq 2$  being  $F$  with mean  $\mu_F$  and the property  $\inf\{x : F(x) = 1\} = \infty$ . Let the associated renewal function be  $m^D$  and  $F$  is not lattice, then for all  $a \geq 0$

$$\lim_{t \rightarrow \infty} m_{t+a}^D - m_t^D = \frac{a}{\mu_F}.$$

If  $F$  and  $G$  are lattice with period  $d$ , then

$$\lim_{n \rightarrow \infty} m_{(n+1)d}^D - m_{nd}^D = \frac{d}{\mu_F}.$$